arXiv:astro-ph/0202144v1 7 Feb 2002

XXX-XXXXX
YYY-YYYYYY
February 1, 2008

Constraining Cosmic Quintessence

G. J. Mathews

Center for Astrophysics, Department of Physics, University of Notre Dame
Notre Dame, IN 46556 USA

K. Ichiki[1,2], T. Kajino[1,2,3] and M. Orito[1]

[1]*National Astronomical Observatory, 2-21-1, Osawa, Mitaka, Tokyo 181-8588, Japan*

[2]*University of Tokyo, Department of Astronomy, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan*

[3]*Graduate University for Advanced Studies, Dept. of Astronomical Science, 2-21-1, Osawa, Mitaka, Tokyo 181-8588, Japan*

and

M. Yahiro

Department of Physics and Earth Sciences, University of the Ryukyus, Nishihara-chou, Okinawa 903-0213, Japan

In quintessence models a scalar field couples with the dominant constituent and only acts like a cosmological constant after the onset of the matter dominated epoch. A generic feature of such solutions, however, is the possibility of significant energy density in the scalar field during the radiation dominated epoch. This possibility is even greater if the quintessence field begins in a kinetic-dominated regime, for example as might be generated at the end of "quintessential inflation." As such, these models can be constrained by primordial nucleosynthesis and the epoch of photon decoupling. Here, we analyze both kinetic dominated and power-law quintessence fields (with and without a supergravity correction). We quantify the allowed initial conditions and effective-potential parameters. We also deduce constraints on the epoch of matter creation at the end of quintessential inflation.

PRESENTED AT

COSMO-01
Rovaniemi, Finland,
August 29 – September 4, 2001

1 Introduction

Observations [1, 2] of Type Ia supernovae and the power spectrum of the cosmic microwave background, together with complementary observational constraints [3], all indicate that the universe is presently accelerating. In one interpretation, this acceleration is due to the influence of a dominant dark energy with a negative pressure. The simplest candidate for such dark energy is a cosmological constant for which the equation of state is $\omega \equiv P/\rho = -1$. A second possibility is derived from the so-called "quintessence" models, in which the dark energy is the result of a scalar field Q slowly evolving along an effective potential $V(Q)$. The equation of state is negative $-1 \leq w_Q \leq 0$, but not necessarily constant.

Introducing a quintessence field helps to reconcile the fine tuning problem and the cosmic coincidence problem associated with a simple cosmological constant. [4]-[11]. Specific forms of the quintessence effective potential can be chosen such that the field Q evolves according to an attractor-like solution. Thus, for a wide variety of initial conditions, the solutions for Q and $\dot{Q}$ rapidly approach a common evolutionary track. These solutions lead naturally to a cross over from an earlier radiation-dominated solution to one in which the universe is dominated by dark energy at late times. Another interesting possible feature is that such models might naturally arise during matter creation at the end of an earlier "quintessential" inflationary epoch [12]. In this case, the Q field emerges in a kinetic-dominated regime at energy densities well above the tracker solution.

It is not yet clear, however, that these models have altogether solved the fine-tuning and cosmic-coincidence problems [13, 14], and there may be some difficulty in embedding quintessence models in string theory [15]. Nevertheless, several such tracker fields have been proposed [10] whose effective potentials may be suggested by particle physics models with dynamical symmetry breaking, by nonperturbative effects [5], by generic kinetic terms "k-essence" in an effective action describing the moduli and massless degrees of freedom in string and supergravity theories [6]-[8], or by static and moving branes in a dilaton gravity background [16].

A general feature of all such solutions, however, is the possibility for a significant contribution of the Q field to the total energy density during the radiation-dominated epoch as well as the present matter-dominated epoch. The yields of primordial nucleosynthesis and the power spectrum of the CMB can be strongly affected by this background energy density. Therefore, we utilize primordial nucleosynthesis and the CMB power spectrum to constrain viable quintessence models.

2 Quintessence Field

A variety of quintessence effective potentials [10] or k-essence effective actions [6]-[8] can be found in the literature. Observational constraints on such quintessence models have been of considerable recent interest [11, 17, 18]. In this paper, we describe the work presented in [11] on Q-field and/or kinetic-dominated quintessence models. We concentrate on an inverse power law for the Q field as originally proposed by Ratra and Peebles [19], $V(Q) = M^{(4+\alpha)}Q^{-\alpha}$, where, M and α are parameters. The parameter M in these potentials is fixed by the condition that $\Omega_Q = 0.7$ at present. Therefore, $\rho_Q(0) = 0.7\rho_c(0) = 5.7h^2 \times 10^{-47}$ GeV^4, and $M \approx (\rho_Q(0)Q^\alpha)^{1/(\alpha+4)}$. If Q is presently near the Planck mass and α is not too small (say $\alpha \gtrsim 4$), this implies a reasonable value [5] for M which resolves the fine tuning problem.

We also consider a modified form of $V(Q)$ as proposed by [20] based upon the condition that the quintessence fields be part of supergravity models. The rewriting of the effective potential in supergravity depends upon choices of the Kähler potential [14]. The flat Kähler potential yields an extra factor of $exp\{3Q^2/2m_{pl}^2\}$ [20]. This comes about by imposing the condition that the expectation value of the superpotential vanishes. The Ratra potential thus becomes $V_{SUGRA}(Q) \rightarrow M^{(4+\alpha)}Q^{-\alpha} \times exp(3Q^2/2m_{pl}^2)$, where the exponential correction becomes largest near the present time as $Q \rightarrow m_{pl}$. This supergravity motivated effective potential is known as the SUGRA potential. The fact that this potential has a minimum for $Q = \sqrt{\alpha/3}m_{pl}$ changes the dynamics. It causes the present value of w_Q to evolve to a cosmological constant ($w_Q \approx -1$) much quicker than for the bare power-law potential [10].

The quintessence field Q obeys the equation of motion $\ddot{Q} + 3H\dot{Q} + dV/dQ = 0$, where the Hubble parameter H is given from the solution to the Friedmann equation, $H^2 = (\dot{a}/a)^2 = 1/m_{pl}^2(\rho_B + \rho_Q)$, where $m_{pl} = (8\pi G/3)^{-1/2} = 4.2 \times 10^{18}$ GeV, ρ_B is the energy density in background radiation and matter, and a is the cosmic scale factor. As the Q field evolves, its contribution to the energy density is given by $\rho_Q = \dot{Q}^2/2 + V(Q)$. Similarly, the pressure in the Q field is $P_Q = \dot{Q}^2/2 - V(Q)$. The equation of state parameter w_Q is a time-dependent quantity, $w_Q \equiv P_Q/\rho_Q = 1 - 2V(Q)/\rho_Q$, where the time dependence derives from the evolution of $V(Q)$ and $\rho(Q)$.

3 Nucleosynthesis Constraint

The quintessence initial conditions are probably set in place near the inflation epoch. By the time of the big bang nucleosynthesis (BBN) epoch, many of the possible initial conditions will have already achieved the tracker solution. However, for initial conditions sufficiently removed from the tracker solution, it is possible that the tracker solution has not yet been achieved by the time of BBN at $0.01 \lesssim T \lesssim 1$ MeV, $10^8 \lesssim z \lesssim 10^{10}$.

For many possible initial conditions the tracker solution is obtained before nucleosynthesis. Along the tracker solution, ρ_Q diminishes in a slightly different way than the radiation-dominant background energy density. For example, as long as $\rho_Q << \rho_B$, the Q-field decays as $\rho_Q \propto a^{-3(1+w_Q)}$, with $w_Q = (\alpha w_B - 2)/(\alpha + 2) < w_B$. The equation of state w_Q is only equal to the background equation of state w_B in the limit $\alpha \rightarrow \infty$. Nevertheless, the tracker solution does not deviate much from ρ_B, even at high redshift for most values of α. Hence, one can characterize the nucleosynthesis results by the (nearly constant) ratio ρ_Q/ρ_B during the BBN epoch. If the energy density in the tracker solution is close to the background energy density, the nucleosynthesis will be affected by the increased expansion rate from the increased total energy density. Such a situation occurs for large values of the power-law exponent α.

A second possibility is that the energy density ρ_Q could exceed the tracker solution and be comparable to or greater than the background energy density during primordial nucleosynthesis. The kinetic energy in the Q field then dominates over the potential energy contribution to ρ_Q and $w_Q = +1$ so that the kinetic energy density diminishes as a^{-6}. In this case there could be a significant contribution from ρ_Q during nucleosynthesis as the Q field approaches the tracker solution. The strongest constraints on this case would arise when ρ_Q is comparable to the background energy density near the time of the weak-reaction freese out, while the later nuclear-reaction epoch might be unaffected. This case is particularly interesting as this kinetic-dominated evolution could be generated by an earlier quintessential inflation epoch [12] as described below.

A final possibility might occur if the Q field approaches the tracker solution from below. In this case, the tracker solution may be achieved after the the BBN epoch so that a small ρ_Q during BBN is easily guaranteed. However, the ultimate tracker curve might still have a large energy density at the later CMB epoch as described below.

Adding energy density from the Q field tends to increase the universal expansion rate. Consequently, the weak reaction rates freeze out at a higher temperature T_w. This fixes the neutron to proton ratio ($n/p \approx \exp[(m_p - m_n)/T_w]$) at a larger value. Since most of the free neutrons are converted into He^4, the primordial helium production is increased. Also, since the epoch of nuclear reactions is shortened, the efficiency of burning deuterium into helium is diminished and the deuterium abundance also increases. Hence, very little excess energy density from the Q field is allowed. The primordial light-element abundances deduced from observations have been reviewed by a number of recent papers [21]-[23]. There are several outstanding uncertainties. For our purposes [11] we adopt $0.226 \le Y_p \le 0.247$ and $2.9 \times 10^{-5} \le D/H \le 4.0 \times 10^{-5}$ as the most relevant constraints.

Figure 1 summarizes allowed values of ρ_Q/ρ_B at $T =1$ MeV ($z \approx 10^{10}$) based upon the nucleosynthesis constraints. The region to the right of the curve labeled D/H corresponds to models in which the primordial deuterium constraint is satisfied, $D/H \le 4.0 \times 10^{-5}$. The region below the line labeled Y_p corresponds to $Y_p \le 0.247$. The hatched region summarizes allowed values for the energy density in the quintessence field during the nucleosynthesis epoch.

In the present work we deduce an absolute upper limit of 5.6% of the background radiation energy density allowed in the quintessence field. This maximum contribution is only allowed for $\eta_{10} \approx 4.75$ or $\Omega_b h^2 \approx 0.017$. A smaller contribution is allowed for other values of η_{10}. Indeed, this optimum η_{10} value is 4σ less than the value implied by the cosmic deuterium abundance [22, 23] $\Omega_b h^2 = 0.020 \pm 0.001$ (1σ) ($\eta_{10} = 5.46 \pm 0.27$). The independent determinations of $\Omega_b h^2$ from high-resolution measurements of the power spectrum of fluctuations in the cosmic microwave background favor a value even higher.



Figure 1: Constraints on the ratio of the energy density in the quintessence field to the background energy density ρ_Q/ρ_B (at $T = 1$ MeV) from the primordial abundances as indicated. The allowed region corresponds to $Y_p \leq 0.247$ and D/H $\leq 4.0 \times 10^{-5}$.

Both the BOOMERANG [26] and DASI [27] data sets imply $\Omega_b h^2 = 0.022^{+0.004}_{-0.003}$ (1σ) $(\eta_{10} = 6.00^{+1.10}_{-0.81})$. The deuterium and CMB constraints together demand that $\eta_{10} \geq 5.19$ which would limit the allowed contribution from the Q field to $\leq 2\%$ of the background energy density.

The most restrictive CMB constraint on $\Omega_b h^2$ derives from demanding a flat universe $(\Omega_{tot} = 1.0)$, and marginalizing the likelihood function over all parameters with assumed Gaussian priors [26] based upon measurements of large-scale structure and Type Ia supernovae. This gives $\Omega_b h^2 = 0.023 \pm 0.003$ (1σ). If one adopts this as a most extreme case, then $\Omega_b h^2 \geq 0.020$. This would correspond to $\eta_{10} \geq 5.46$. From Figure 1 this would imply a much more stringent constraint that only about 0.1% of the background energy density could be contributed by the Q field. Of course, this is only a 1σ constraint, and the upper limit to Y_p is not well enough established to rule out a contribution to the energy density at the 0.1% level. We adopt the conservative constraint of 5.6%. Nevertheless, it is of interest to explore how the quintessence parameters allowed by BBN might improve should the constraints from BBN ever be so tightly defined. Therefore, we will consider 0.1% as a conceivable limit that demonstrates the sensitivity to BBN. Even the most conservative 5.6% limit adopted here corresponds to only about half of the energy density allowed in [24] for 3 neutrino flavors.

4 Equation of State Constraint

The Q field must behave like dark energy during the present matter-dominated epoch, i.e. the equation of state should be sufficiently negative, ($w_Q \equiv P_Q/\rho_Q < 0$) by the present time. We adopt observational constraints on w_Q from [3]. They adopted a most conservative approach based upon progressively less reliable data sets. Using the most reliable current low-redshift and microwave background measurements, the 2σ limits are $-1 < w_Q < -0.2$. Factoring in the constraint from Type Ia supernovae reduces the range to $-1 < w_Q < -0.4$. This derives from a concordance analysis of models consistent with each observational constraint at the 2σ level or better. A combined maximum likelihood analysis suggests a smaller range of $-0.8 < w_Q < -0.6$ for quintessence models which follow the tracker solution, though $w_Q \approx -1$ is still allowed in models with nearly a constant dark energy. We invoke these three possible limit ranges for the present value of w_Q.

5 CMB Constraint

There are two effects on the epoch of photon decoupling to be considered. In the case where the energy density in the quintessence field is negligible during photon decoupling [10] the only effect of the dark energy is to modify the angular distance-redshift relation [25]. The existence of dark energy along the look-back to the surface of last scattering shifts the acoustic peaks in the CMB power spectrum to smaller angular scales and larger l-values. The amplitude of the first acoustic peak in the power spectrum also increases, but not as much for quintessence models as for a true cosmological constant Λ. The basic features of the observed power spectrum [26] can be fit [10] with either of the quintessence potentials considered here. For our purposes, this look-back constraint is already satisfied by demanding that $\Omega_Q = 0.7$ at the present time.

The second effect occurs if the energy density in the Q field is a significant fraction of the background energy during the epoch of photon decoupling. Then it can increase the expansion rate and push the l values for the acoustic peaks to larger values and increase their amplitude [25]. Such an effect has been considered by a number of authors in various contexts [17, 25, 28]. For our purposes, we adopt a constraint [17] based upon the latest CMB sky maps of the Boomerang [26] and DASI [27] collaborations. The density in the Q field can not exceed $\Omega_Q \leq 0.39$ during the epoch of photon decoupling. This implies a maximum of $\rho_Q/\rho_B = \Omega_Q/(1-\Omega_Q) \lesssim 0.64$ during photon decoupling.

6 Quintessential Inflation

Another possible constraint arises if the kinetic term dominates at an early stage. In this case $\rho_Q \approx \dot{Q}^2/2$ and ρ_Q decreases with scale factor as a^{-6}. At very early times this kinetic regime can be produced by so-called "quintessential inflation" [12]. In this paradigm entropy in the matter fields comes from gravitational particle production at the end of inflation. The universe is presumed to exit from inflation directly into a kinetic-dominated quintessence regime during which the matter is generated. An unavoidable consequence of this process, however, is the generation of gravitational waves along with matter and the quanta of the quintessence field [12, 29, 30, 31] at the end of inflation.

6.1 Energy in Quanta and Gravity Waves

The energy density in created particles can be deduced using quantum field theory in curved space-time [12, 30, 31], $\rho_B \approx (1/128) N_s H_1^4 (z + 1/z_1 + 1)^4 (g_1/g_{eff}(z))^{1/3}$, where H_1 and z_1 are the expansion factor and redshift at the matter thermalization epoch at the end of quintessential inflation, respectively. The factor of 128 in this expression comes from the explicit integration of the particle creation terms.

When the gravitons and quanta of the Q field are formed at the end of inflation, one expects [12] the energy density in gravity waves to be twice the energy density in the Q-field quanta (because there are two graviton polarization states). In this paradigm then, wherever we have deduced a constraint on ρ_Q, it should be taken as the sum of three different contributions. One is the dark energy from the vacuum expectation value $\langle \rho_Q \rangle$ of the Q field; a second is from the fluctuating part $\rho_{\delta Q}$ of the Q field; and a third is from the energy density ρ_{GW} in relic gravity waves. Thus, we have $\rho_Q \rightarrow (\langle \rho_Q \rangle + \rho_{\delta Q} + \rho_{GW})$. The energy density in gravity waves and quanta scales like radiation after inflation, $\rho_{GW} + \rho_{\delta Q} \propto a^{-4}$, while the quintessence field vacuum expectation value evolves as $\langle \rho_Q \rangle \propto a^{-6}$ during the kinetic dominated epoch. This epoch following inflation lasts until the energy in the Q field falls below the energy in background radiation and matter, $\rho_Q \leq \rho_B$.

Thus, for the kinetic dominated initial conditions, gravity waves could be an important contributor to the excess energy density during nucleosynthesis. The relative contribution of gravity waves and quintessence quanta compared to the background matter fields is just given by the relative number of degrees of freedom. At the end of inflation, the relative fraction of energy density in quanta and gravity waves is given by [12] $(\rho_{\delta Q} + \rho_{GW})/\rho_B = 3/N_s$, where N_s is the number of ordinary scalar fields at the end of inflation. In the minimal supersymmetric model $N_s = 104$. Propagating this ratio to the time of nucleosynthesis requires another factor of $(g_n/g_1)^{1/3}$ where $g_n = 10.75$ counts the number of effective relativistic degrees of freedom just before electron-positron annihilation, and g_1 counts the number of degrees of freedom during matter thermalization after the end of inflation. In the minimal standard model $g_1 = 106.75$, but in supergravity models this increases to $\sim 10^3$.

Combining these factors we have $(\rho_{\delta Q} + \rho_{GW})/\rho_B \leq 0.014$, during nucleosynthesis. Hence, in this paradigm, the allowed values of ρ_Q/ρ_B consistent with nucleosynthesis could be reduced from a maximum of 0.056 to 0.042, further tightening the constraints deduced here.

6.2 Gravity-Wave Spectrum

There has been considerable recent interest [29, 30] in the spectrum of gravity waves produced in the quintessential inflation paradigm. One might expect that the *COBE* constraints on the spectrum also lead to constraints on the Q field. However, we conclude below that no significant constraint on the initial ρ_Q or effective potential is derived from the gravity wave spectrum. On the other hand, the BBN and CMB gravity-wave constraints can be used to provide useful constraints on the quintessential inflation epoch as we now describe. Our argument is as follows: The logarithmic gravity-wave energy spectrum observed at the present time can be defined in terms of a differential closure fraction, $\Omega_{GW}(\nu) \equiv (1/\rho_c)(d\rho_{GW}/d\ln\nu)$, where the ρ_{GW} is the present energy density in relic gravitons and $\rho_c(0) = 3H_0^2/8\pi G = H_0^2 m_{pl}^2$ is the critical density. This spectrum has been derived in several recent papers [29, 30]. It is characterized by spikes at low and high frequency. The most stringent constraint at present derives from the *COBE* limit on the tensor component of the CMB power spectrum at low multipoles. There is also a weak constraint from the binary pulsar [30] and an integral constraint from nucleosynthesis as mentioned above.

For our purposes, the only possible new constraint comes from the *COBE* limits on the tensor component of the CMB power spectrum. The soft branch in the gravity-wave spectrum lies in the frequency range between the present horizon $\nu_0 = 1.1\times 10^{-18}\Omega_M^{1/2} h$ Hz and the decoupling frequency $\nu_{dec}(0) = 1.65\times 10^{-16}\Omega_M^{1/2} h$ Hz, where we adopt $\Omega_M = 0.3$ for the present matter closure fraction. The constraint on the spectrum can be written [30],

$$\Omega_{GW}(\nu) = \Omega_\gamma \frac{81}{(16\pi)^2}\left(\frac{g_{dec}}{g_1}\right)^{1/3}\left(\frac{H_1}{m_{pl}}\right)^2 \times \left(\frac{\nu_{dec}}{\nu}\right)^2 \times \ln^2\left(\frac{\nu_r}{\nu_1}\right) \leq 6.9h^{-2}\times 10^{-11} \quad , \tag{1}$$

where, $\Omega_\gamma = 2.6\times 10^{-5}h^2$ is the present closure fraction in photons. The number of relativistic degrees of freedom at decoupling is $g_{dec} = 3.36$. As noted previously, g_1 is the number of relativistic degrees of freedom after matter thermalization. In the minimal standard model is $g_1 = 106.75$. The quantity H_1 is the expansion rate at the end of inflation. It is simply related to the kinetic energy ρ_Q after inflation, $\rho_Q(z_1) = H_1^2 m_{pl}^2$. The logarithmic term in Eq. (1) involves the ratio of the present values of the frequency $\nu_r(0)$ characteristic of the start of radiation domination ($\rho_B = \rho_Q$ at $z = z_r$), to the frequency characteristic of matter thermalization at the end of inflation $\nu_1(0)$. This ratio is just $\nu_r(0)/\nu_1(0) = (z_r + 1/z_1 + 1)^2$. The identity $\rho_B = \rho_Q$ at $z = z_r$ then gives, $\nu_r/\nu_1 = N_s/128(H_1/m_{pl})^2(g_1/g_r)^{1/3}$, where for the cases of interest $g_r = 10.75$.

Collecting these terms, we can then use Eq. (1) to deduce a constraint on the expansion factor at the end of inflation $H_1^2 < 1.4\times 10^{-11} m_{pl}^2$. For kinetic dominated models, $\rho_Q(z_1)$ at the end of inflation is simply related to the energy density ρ_Q at z, $\rho_Q(z) = \rho_Q(z_1)[(z+1)/(z_1+1)]^6$. Similarly the background matter energy density scales as $\rho_B(z) = \rho_B(z_1)(g_1/g_{eff}(z))^{1/3}[(z+1)/(z_1+1)]^4$.

Considering the present energy density in photons and neutrinos, we can find a relation between H_1 and z_1: $\rho_{\gamma 0} + \rho_{\nu 0} = 1.1 \times 10^{-125} m_{pl}^4 = (N_s/128) H_1^4 (g_1/g_{dec})^{1/3} (z_1+1)^{-4}$. We then deduce that $z_1 < 8.4 \times 10^{25}$ and there is only a lower limit on ρ_Q/ρ_B given by the constraint on H_1. At our initial epoch $z = 10^{12}$, we find, $\rho_Q(z)/\rho_B(z) \gtrsim 5.6 \times 10^{-18}$. This limit is not particularly useful because ρ_Q field must exceed ρ_B at z_1 in order for the gravitational particle production paradigm to work. The implication is then that all initial conditions in which the kinetic term dominates over the background energy at z_1 are allowed in the quintessential inflation scenario. Hence, we conclude that the gravity-wave spectrum does not presently constrain the initial ρ_Q or $V(Q)$ in the quintessential inflation model.

However, the limits on $\rho_Q/\rho_B \leq 560$ derived from the BBN constraints discussed below can be used to place a lower limit on the expansion rate at the end of inflation in this model. This in turn implies a lower limit on the redshift for the end of quintessential inflation. Thus, we have $2.2 \times 10^{-21} < H_1^2/m_{pl}^2 < 1.45 \times 10^{-11}$, and $1.0 \times 10^{21} < z_1 < 8.4 \times 10^{25}$ in the minimal supersymmetric model.

This implies that quintessential inflation must end at an energy scale somewhere between about 10^8 and 10^{13} GeV, well below the Planck scale. By similar reasoning one can apply this argument to the gravity-wave spectrum from normal inflation as given in [30]. We deduce an upper limit to the epoch of matter thermalization of $H_1 \leq 3.1 \times 10^{-10} m_{pl}^2$ which implies $z_1 \leq 7.3 \times 10^{28} [g(z_1)/3.36]^{1/12}$. In this case there is no lower limit from BBN as there is no Q field present after inflation.

7 Results and Discussion

The equations of motion were evolved for a variety of initial Q field strengths and power-law parameters α. As initial conditions, the quintessence field was assumed to begin with equipartition, i.e. $\dot{Q}^2/2 = V(Q)$, and $w_Q = 0$. This seems a natural and not particularly restrictive choice, since w_Q quickly evolves toward the kinetic ($w_Q = +1$) or the tracker solution depending upon whether one begins above or below the tracker curve.

Constraints on α and the initial value for the Q-field energy density $\rho_Q(z)/\rho_B(z)$ at $z = 10^{12}$ were deduced numerically. These are summarized in Figure 2 for both: (a) the bare Ratra power-law potential; and (b) its SUGRA corrected form. Our constraints can easily be converted to closure fraction by $\Omega_Q = (\rho_Q/\rho_B)/(1 + \rho_Q/\rho_B)$. For purposes of illustration, we have arbitrarily specified initial conditions at $z = 10^{12}$, corresponding to $T \sim 10^{12}$ K, roughly just after the time of the QCD epoch. At any time the energy density $\rho_{rel}(z)$ in relativistic particles is just $\rho_{rel}(z) = \rho_{\gamma 0}(3.36/g_{eff}(z))^{1/3}(z+1)^4$, where $\rho_{\gamma 0} = 2.0 \times 10^{-51}$ GeV^4, is the present energy density in microwave background photons, and we take $g_{eff}(z) = 10.75$ between $z = 10^{12}$ and the beginning of BBN just before electron-positron annihilation ($z \approx 10^{10}$).

The envelope of models which obtain a tracker solution by the epoch of nucleosynthesis



Figure 2: Contours of allowed values for α and initial ρ_Q/ρ_B (at $z = 10^{12}$) from various constraints as indicated for (a) the bare power-law potential, and (b) the SUGRA corrected potential. Models in which the tracker solution is obtained by the BBN epoch are indicated by the upper and lower curves. Values of α to the right of the lines labeled $w_Q = -0.6, -0.4, -0.2$ on (a) are excluded by the requirement that the present equation of state be sufficiently negative. The BBN constraint for a maximum energy density in the Q field of 0.1% (dotted line) and 5.6% (solid line) are also indicated. For the SUGRA potential (b) all tracker solutions to the right of the region labeled $w_Q = -.8$ are allowed.

are indicated by upper and lower curved lines in Figures 2a and 2b. The general features of these constraints are as follows. If the initial energy density in the Q field is too large, the tracker solution is not reached by the time of BBN. The Q-field energy density can then significantly exceed the background energy during nucleosynthesis. This situation corresponds to the excluded regions on the top of Figures 2a and 2b. All solutions consistent with the primordial nucleosynthesis constraints are also consistent with our adopted CMB Q-energy constraint as also shown at the top of Figures 2a and 2b.

The excluded regions at the top and bottom of figures 2a and 2b can be easily understood analytically. For example, the excluded (no Λ) region at the bottom of these figures reflects the fact that if ρ_Q is initially below the value presently required by $\Omega_\Lambda = 0.7$ it can not evolve toward a larger value. Hence, $\rho_Q/\rho_B < 2.8 \times 10^{-44}$ is ruled out for $h = 0.7$. Similarly, the "Excluded by BBN" region comes from requiring that $\rho_Q(z_{BBN})/\rho_B(z_{BBN}) < 0.056$. For this constraint we are only considering cases in which the Q field is approaching the tracker solution from above during nucleosynthesis. Hence, it is in the kinetic regime in which $\rho_Q \propto z^6$ while the background scales as z^4. Thus, we have $\rho_Q(z=10^{12})/\rho_B(z=10^{12}) > 0.056(10^{12}/10^{10})^{6-4} = 560$ is excluded. By similar reasoning, the "Excluded by CMB" region is given by $\rho_Q(z_{CMB})/\rho_B(z_{CMB}) > 0.64$, or $\rho_Q(z=10^{12})/\rho_B(z=10^{12}) > 0.64(10.75/3.36)^{1/3}(10^{12}/10^3)^{6-4} = 9.4 \times 10^{17}$.

For the bare Ratra power-law potential (Fig. 2a) the main constraint is simply the requirement that the equation of state be sufficiently negative by the present time. The sensitivity of the allowed power-law exponent to the equation of state is indicated by the w_Q = -0.2, -0.4, and -0.6 lines on Figure 2a. In the present Q-dominated epoch, lines of constant w_Q must be evaluated numerically. The slight slope to these curves comes from the fact that $V(Q)/\rho(Q)$ has not yet reached unity, i.e. there is still some small kinetic contribution to $\rho(Q)$ and the amount of kinetic contribution depends upon α.

For the bare Ratra power-law potential, tracker solutions with $\alpha \lesssim 20$ are allowed if $w_Q \leq -0.2$. The allowed values for α reduce to < 9 and < 2 if the more stringent -0.4 and -0.6 constraints are adopted. However, if α is too small, say $\alpha \leq 2$, then the potential parameter M becomes a very small fraction of the Planck mass and the fine tuning problem is reintroduced.

For models in which the tracker solution is obtained by the time of BBN, the potential parameters are only constrained if the most conservative equation of state limit ($w_Q < -0.2$) and most stringent nucleosynthesis constraint ($\rho_Q/\rho_B < 0.1\%$) are adopted. On the other hand, independently of the equation of state constraint, nucleosynthesis limits a large family of possible kinetic-dominated solutions even though they provide the correct present dark energy.

For the SUGRA-corrected Q fields (Fig. 2b), the constraint from w_Q is greatly relaxed. In fact, w_Q is sufficiently negative ($w_Q < -0.6$) for all $\alpha < 10^4$. The reason is that all tracker solutions have $w_Q \approx -1$. This is because w_Q decays much faster toward -1 for the SUGRA potential. Also, the potential has a finite minimum which is equal to the present dark-energy density. The Q field quickly evolves to near the potential minimum and has negligible kinetic energy by the present time. Any potential which becomes flat at late times gives $w_Q \approx -1$ and the dark energy looks like a cosmological constant. All SUGRA models which achieve the tracker solution also have a small ρ_Q during primordial nucleosynthesis. Hence, there is no constraint from nucleosynthesis except for those kinetic-dominated models in which the Q field is still far above the tracker solution during the nucleosynthesis epoch.

We do note, however, that if a lower limit of $w_Q > -0.8$ is adopted for tracker solutions from [9], then only a power law with $\alpha \gtrsim 30$ is allowed. This makes the SUGRA potential the preferred candidate for quintessence. The large α implies values of M close to the Planck mass, thus avoiding any fine tuning problem. However, this potential will be constrainable by BBN if the light-element constraints become sufficiently precise to limit ρ_Q at the 0.1% level.

8 Conclusions

We conclude that for both the bare Rata inverse power-law potential and its SUGRA-corrected form, the main constraints for models which achieve the tracker solution by the nucleosynthesis epoch is from the requirement that the equation of state parameter becomes sufficiently negative by the present epoch. The main constraint from nucleosynthesis is for models which are kinetic dominated ar the time of nucleosynthesis. The SUGRA-corrected potential is the least constrained and avoids the fine tuning problem for M. Therefore, it may be the preferred candidate for the quintessence field, although BBN may eventually limit this possibility. We also note that the constraints considered here provide useful constraints on the regime of matter creation at the end of quintessential inflation.

Work supported in part by the Grant-in-Aid for Scientific Research (10640236, 10044103, 11127220, 12047233) of the Ministry of Education, Science, Sports, and Culture of Japan, and also by DoE Nuclear Theory Grant (DE-FG02-95-ER40934 at UND).

References

[1] P. M. Garnavich, et al., Astrophys. J., **509**, (1998) 74; Riess, et al., Astron. J, **116**, (1998) 1009.

[2] S. Perlmutter, et al., Nature **391**, (1998) 51; Astrophys. J., **517**, (1998) 565.

[3] L. Wang, R.R. Caldwell, J.P. Ostriker and P. J. Steinhardt, Astrophys. J., 530, (2000) 17.

[4] C. Wetterich, Nucl. Phys., B302, (1988) 668.

[5] I. Zlatev, L. Wang, and P. J. Steinhardt, Phys. Rev. Lett., 82, (1999) 896.

[6] T. Chiba, T. Okabe, and M. Yamaguchi, Phys. Rev. D62, (2000) 023511.

[7] C. Armendariz-Picon, V. Mukhanov, and P. J. Steinhardt, Phys. Rev Lett., **85**, (2000) 4438.

[8] C. Armendariz-Picon, V. Mukhanov, and P. J. Steinhardt, Phys. Rev D63, (2001) 103510.

[9] P. J. Steinhardt, L. Wang and I. Zlatev, Phys. Rev. D59, (1999) 123504.

[10] P. Brax, J. Martin, and A. Riazuelo, Phys. Rev D62, (2000) 103505.

[11] M. Yahiro, G. J. Mathews, K. Ichiki, T. Kajino, Phys. Rev. D, (2002) in press.

[12] P.J.E. Peebles and A. Vilenkin, Phys. Rev. D59, (1999) 063505.

[13] C. Kolda and D. H. Lyth, Phys. Lett., **B458**, (1999) 197; S. Weinberg, in Proc. Dark Matter 2000, Marina Del Rey, Feb, 2000, astro-ph/0005265, (2000).

[14] E. J. Copeland, N. J. Nunes, and F. Rosati, Phys. Rev. D62, (2000) 123503.

[15] T. Banks, JHEP submitted (2000), hep-th/0007146; T. Banks and W. Fischler, JHEP submitted (2000), hep-th/0102077; S. Hellerman, N. Kaloper, and L. Susskind, JHEP, 06, 003 (2001); W. Fischler et al., JHEP, 07, 003 (2001) hep-th/0104181; C. Kolda and W. Lahneman, hep-th/0105300.

[16] B. Chen and F-L. Lin, Phys. Rev. D65, (2002) 044007.

[17] R. Bean, S. H. Hansen, and A. Melchiorri, Phys. Rev. D64, (2001) 103508.

[18] X. Chen, R. J. Scherrer, and G. Steigman, Phys. Rev D63, (2001) 123504.

[19] B. Ratra and P.J.E. Peebles, Phys. Rev. D37, (1988) 3406; P.J.E. Peebles and B. Ratra, Astrophys. J. Lett., 325, (1988) L17.

[20] P. Brax and J. Martin, Phys. Lett., **B468**, (1999) 40; Phys.Rev. D61 (2000) 103502; P. Brax, J. Martin, A. Riazuelo, Phys. Rev. **D62**, (2000) 103505.

[21] K.A. Olive, G. Steigman, and T.P.Walker, Phys. Rep. 333, (2000) 389.

[22] K.M. Nollett and S. Burles, Phys. Rev. **D61**, (2000) 123505; S. Burles, K.M. Nollett, & M.S. Turner, Phys. Rev. **D63**, (2001) 063512.

[23] D. Tytler, J. M. O'Meara, N. Suzuki, & D. Lubin Phys. Rep., 333, (2000) 409.

[24] K. Freese, F. Adams, J. Frieman, and E. Mottola, in *Origin and Distribution of the Elements*, G. J. Mathews, ed. (World Scientific: Singapore) (1988), pp. 97-115.

[25] W. Hu, D. Scott, N. Sugiyama, and M. White, Phys. Rev. D52, (1995) 5498.

[26] C.B. Netterfield, et al. (Boomerang Collaboration), Submitted to Astrophys. J. (2001), astro-ph/0104460.

[27] C. Pryke, et al. (DASI Collaboration), Submitted to Astrophys. J. (2001), astro-ph/0104490.

[28] S. Hannestad, Phys. Rev. Lett., 85, (2000) 4203.

[29] A. Riazuelo, and J.-P. Uzan, Phys. Rev. **D62**, (2000) 083506.

[30] M. Giovannini, Class. Quant. Grav **16**, (1999) 2905; Phys. Rev. D60, (1999) 123511.

[31] L. H. Ford, Phys. Rev. **D35**, (1987) 2955.